

Supplement of Biogeosciences, 16, 329–346, 2019
<https://doi.org/10.5194/bg-16-329-2019-supplement>
© Author(s) 2019. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

On the role of climate modes in modulating the air–sea CO₂ fluxes in eastern boundary upwelling systems

Riley X. Brady et al.

Correspondence to: Riley X. Brady (riley.brady@colorado.edu)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

California Current NPGO Composites





Figure S1: CalCS composites for the positive and negative phase of the NPGO for (a) sea surface temperature, (b) wind stress and wind stress curl, and (d) 50m dissolved inorganic carbon anomalies. The black outline in these plots shows our statistical study region. The positive NPGO forcing pattern for the North Pacific is displayed in (c) for wind stress and sea surface height anomalies. The gray box in (c) delineates the region displayed in (a), (b), and (c). Composites were calculated by averaging every NPGO event that exceeded two standard deviations above or below the index mean for the full ensemble.

California Current Spatial CO₂ Flux Decomposition

(a) California Current 1 σ NPGO Ensemble Mean Response



(b) California Current 1 σ PDO Ensemble Mean Response




Figure S2: Linear Taylor expansion by grid cell for CO₂ flux anomalies in the CalCS in response to (a) a 1 σ NPGO and (b) a 1 σ PDO.

California Current PDO Composites




Figure S3: As in Figure S1, but for the PDO.

Humboldt Current Nino3 Composites




Figure S4: As in Figure S1, but for the HumCS response to ENSO.

Canary Current NAO Composites




Figure S5: As in Figure S1, but for the CanCS response to the NAO.

Humboldt and Canary Current Spatial CO₂ Flux Decomposition

(a) Humboldt Current 1° Nino3 Ensemble Mean Response




Figure S6: As in Figure S2, but for (a) the HumCS response to ENSO, and (b) the CanCS response to the NAO.

Relative Contributions to sDIC Tendency Anomaly by Grid Cell




Figure S7: Grid cell analysis of the relative contributions to the sDIC tendency anomaly for (a) the CalCS response to the NPGO, (b) the CalCS response to the PDO, (c) the HumCS response to ENSO, and (d) the CanCS response to the NAO. Each subplot contains (from left to right) the relative contribution of CO₂ flux, circulation, and biology anomalies.